

Search for CP Violation in the Decays $D^0 \rightarrow K_S^0 P^0$

B. R. Ko,¹⁷ E. Won,¹⁷ K. Arinstein,^{1,33} T. Aushev,^{19,12} A. M. Bakich,⁴⁰ V. Balagura,¹² E. Barberio,²³ K. Belous,¹¹ V. Bhardwaj,³⁵ B. Bhuyan,⁹ M. Bischofberger,²⁵ A. Bondar,^{1,33} A. Bozek,²⁹ M. Bračko,^{21,13} J. Brodzicka,²⁹ T. E. Browder,⁷ A. Chen,²⁶ P. Chen,²⁸ B. G. Cheon,⁶ C.-C. Chiang,²⁸ I.-S. Cho,⁵⁰ K. Cho,¹⁶ K.-S. Choi,⁵⁰ S.-K. Choi,⁵ Y. Choi,³⁹ S. Eidelman,^{1,33} D. Epifanov,^{1,33} M. Feindt,¹⁵ N. Gabyshev,^{1,33} A. Garmash,^{1,33} B. Golob,^{20,13} H. Ha,¹⁷ J. Haba,⁸ H. Hayashii,²⁵ Y. Horii,⁴⁵ Y. Hoshi,⁴⁴ W.-S. Hou,²⁸ H. J. Hyun,¹⁸ T. Iijima,²⁴ K. Inami,²⁴ A. Ishikawa,³⁶ R. Itoh,⁸ M. Iwabuchi,⁵⁰ T. Iwashita,²⁵ T. Julius,²³ J. H. Kang,⁵⁰ T. Kawasaki,³¹ C. Kiesling,²² H. O. Kim,¹⁸ M. J. Kim,¹⁸ Y. J. Kim,¹⁶ K. Kinoshita,³ P. Kodyš,² S. Korpar,^{21,13} P. Križan,^{20,13} R. Kumar,³⁵ A. Kuzmin,^{1,33} Y.-J. Kwon,⁵⁰ S.-H. Kyeong,⁵⁰ M. J. Lee,³⁸ S.-H. Lee,¹⁷ C. Liu,³⁷ D. Liventsev,¹² R. Louvot,¹⁹ A. Matyja,²⁹ K. Miyabayashi,²⁵ H. Miyata,³¹ Y. Miyazaki,²⁴ R. Mizuk,¹² G. B. Mohanty,⁴¹ T. Mori,²⁴ E. Nakano,³⁴ M. Nakao,⁸ S. Nishida,⁸ K. Nishimura,⁷ O. Nitoh,⁴⁸ S. Ogawa,⁴³ T. Ohshima,²⁴ S. Okuno,¹⁴ S. L. Olsen,^{38,7} P. Pakhlov,¹² C. W. Park,³⁹ H. Park,¹⁸ H. K. Park,¹⁸ R. Pestotnik,¹³ M. Petrič,¹³ L. E. Pilonen,⁴⁹ A. Poluektov,^{1,33} M. Röhrken,¹⁵ Y. Sakai,⁸ O. Schneider,¹⁹ C. Schwanda,¹⁰ A. J. Schwartz,³ K. Senyo,²⁴ M. E. Sevier,²³ M. Shapkin,¹¹ V. Shebalin,^{1,33} C. P. Shen,⁷ J.-G. Shiu,²⁸ B. Shwartz,^{1,33} F. Simon,^{22,42} J. B. Singh,³⁵ P. Smerkol,¹³ Y.-S. Sohn,⁵⁰ E. Solovieva,¹² S. Stanič,³² M. Starič,¹³ M. Sumihama,^{51,4} K. Sumisawa,⁸ T. Sumiyoshi,⁴⁷ S. Tanaka,⁸ Y. Teramoto,³⁴ K. Trabelsi,⁸ M. Uchida,^{51,46} S. Uehara,⁸ T. Uglov,¹² Y. Unno,⁶ Y. Usov,^{1,33} G. Varner,⁷ K. E. Varvell,⁴⁰ A. Vinokurova,^{1,33} C. H. Wang,²⁷ M.-Z. Wang,²⁸ Y. Watanabe,¹⁴ Y. Yamashita,³⁰ M. Yamauchi,⁸ Z. P. Zhang,³⁷ V. Zhilich,^{1,33} V. Zhulanov,^{1,33} A. Zupanc,¹⁵ and O. Zyukova,^{1,33}

(The Belle Collaboration)

¹*Budker Institute of Nuclear Physics, Novosibirsk*

²*Faculty of Mathematics and Physics, Charles University, Prague*

³*University of Cincinnati, Cincinnati, Ohio 45221*

⁴*Gifu University, Gifu*

⁵*Gyeongsang National University, Chinju*

⁶*Hanyang University, Seoul*

⁷*University of Hawaii, Honolulu, Hawaii 96822*

⁸*High Energy Accelerator Research Organization (KEK), Tsukuba*

⁹*Indian Institute of Technology Guwahati, Guwahati*

¹⁰*Institute of High Energy Physics, Vienna*

¹¹*Institute of High Energy Physics, Protvino*

¹²*Institute for Theoretical and Experimental Physics, Moscow*

¹³*J. Stefan Institute, Ljubljana*

¹⁴*Kanagawa University, Yokohama*

¹⁵*Institut für Experimentelle Kernphysik, Karlsruher Institut für Technologie, Karlsruhe*

¹⁶*Korea Institute of Science and Technology Information, Daejeon*

¹⁷*Korea University, Seoul*

¹⁸*Kyungpook National University, Taegu*

¹⁹*École Polytechnique Fédérale de Lausanne (EPFL), Lausanne*

²⁰*Faculty of Mathematics and Physics, University of Ljubljana, Ljubljana*

²¹*University of Maribor, Maribor*

²²*Max-Planck-Institut für Physik, München*

²³*University of Melbourne, School of Physics, Victoria 3010*

²⁴*Nagoya University, Nagoya*

²⁵*Nara Women's University, Nara*

²⁶*National Central University, Chung-li*

²⁷*National United University, Miao Li*

²⁸*Department of Physics, National Taiwan University, Taipei*

²⁹*H. Niewodniczanski Institute of Nuclear Physics, Krakow*

³⁰*Nippon Dental University, Niigata*

³¹*Niigata University, Niigata*

³²*University of Nova Gorica, Nova Gorica*

³³*Novosibirsk State University, Novosibirsk*

³⁴*Osaka City University, Osaka*

³⁵*Panjab University, Chandigarh*

- ³⁶Saga University, Saga
³⁷University of Science and Technology of China, Hefei
³⁸Seoul National University, Seoul
³⁹Sungkyunkwan University, Suwon
⁴⁰School of Physics, University of Sydney, NSW 2006
⁴¹Tata Institute of Fundamental Research, Mumbai
⁴²Excellence Cluster Universe, Technische Universität München, Garching
⁴³Toho University, Funabashi
⁴⁴Tohoku Gakuin University, Tagajo
⁴⁵Tohoku University, Sendai
⁴⁶Tokyo Institute of Technology, Tokyo
⁴⁷Tokyo Metropolitan University, Tokyo
⁴⁸Tokyo University of Agriculture and Technology, Tokyo
⁴⁹CNP, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061
⁵⁰Yonsei University, Seoul
⁵¹Research Center for Nuclear Physics, Osaka

We have searched for CP violation in the decays $D^0 \rightarrow K_S^0 P^0$ where P^0 denotes a neutral pseudo-scalar meson that is either a π^0 , η , or η' using KEKB asymmetric-energy e^+e^- collision data corresponding to an integrated luminosity of 791 fb^{-1} collected with the Belle detector. No evidence of significant CP violation is observed. We report the most precise CP asymmetry measurement in the decay $D^0 \rightarrow K_S^0 \pi^0$ to date: $A_{CP}^{D^0 \rightarrow K_S^0 \pi^0} = (-0.28 \pm 0.19 \pm 0.10)\%$. We also report the first measurements of CP asymmetries in the decays $D^0 \rightarrow K_S^0 \eta$ and $D^0 \rightarrow K_S^0 \eta'$: $A_{CP}^{D^0 \rightarrow K_S^0 \eta} = (+0.54 \pm 0.51 \pm 0.16)\%$ and $A_{CP}^{D^0 \rightarrow K_S^0 \eta'} = (+0.98 \pm 0.67 \pm 0.14)\%$, respectively.

PACS numbers: 11.30.Er, 13.25.Ft, 14.40.Lb

The recent evidence for $D^0 - \bar{D}^0$ mixing [1–3] and the corresponding mixing parameters [4] are at the upper edge of standard model (SM) predictions [5]. However, large theoretical uncertainties in these predictions limit the sensitivity to effects of physics beyond the SM. An alternative, potentially more promising approach to search for new physics (NP) is the study of violation of the combined Charge-conjugation and Parity symmetries (CP) in the decays of charmed mesons [6]. In contrast to mixing, the expected SM CP violation in the charm sector is small [7].

In this Letter we report time-integrated CP asymmetry measurements in the decays $D^0 \rightarrow K_S^0 P^0$ [8] where P^0 denotes a neutral pseudo-scalar meson: π^0 , η , or η' . The time-integrated asymmetry, A_{CP} , is defined as

$$A_{CP}^{D^0 \rightarrow K_S^0 P^0} = \frac{\Gamma(D^0 \rightarrow K_S^0 P^0) - \Gamma(\bar{D}^0 \rightarrow K_S^0 P^0)}{\Gamma(D^0 \rightarrow K_S^0 P^0) + \Gamma(\bar{D}^0 \rightarrow K_S^0 P^0)}, \quad (1)$$

where Γ is the partial decay width.

The observed $K_S^0 P^0$ final states are mixtures of $D^0 \rightarrow \bar{K}^0 P^0$ and $D^0 \rightarrow K^0 P^0$ decays where the former are Cabibbo-favored (CF) and the latter are doubly Cabibbo-suppressed (DCS). In the absence of direct CP violation in CF and DCS decays, as expected in the SM, the CP violation in these processes within the SM is generated from mixing and interference of decays with and without mixing, which is parameterized by a^{ind} (we adopt the symbols used in Ref. [6]). SM $K^0 - \bar{K}^0$ mixing leads to a small CP asymmetry in final states containing a neutral kaon, even if no CP violating phase exists in the charm decay. The asymmetry that is expected

from the SM is measured to be $(-0.332 \pm 0.006)\%$ [9] from K_L^0 semileptonic decays and referred to as $A_{CP}^{\bar{K}^0}$ [10], which is reflected in the value of $A_{CP}^{D^0 \rightarrow K_S^0 P^0}$ if DCS decay contributions are ignored. Since the a^{ind} value expected from the SM is at most $\mathcal{O}(10^{-4})$ [6, 7], the value of CP asymmetry in the decays $D^0 \rightarrow K_S^0 P^0$ within the SM is approximately $A_{CP}^{\bar{K}^0}$. On the other hand, if NP processes contain additional weak phases other than the one in the Kobayashi-Maskawa ansatz [11], interferences between CF and DCS decays could generate $\mathcal{O}(1)\%$ direct CP asymmetry in the decays $D^0 \rightarrow K_S^0 P^0$ [12]. Physics beyond the SM could also induce $\mathcal{O}(1)\%$ indirect CP asymmetry [6]. Thus, observing A_{CP} inconsistent with $A_{CP}^{\bar{K}^0}$ in $D^0 \rightarrow K_S^0 P^0$ decays would be strong evidence for processes involving physics beyond the SM [6, 12].

In addition to A_{CP} measurements, we examine the universality of a^{ind} in D^0 decays [6] by comparing our previous result [2] with the $A_{CP}^{D^0 \rightarrow K_S^0 \pi^0}$ value reported in this Letter. Our previously measured values of direct CP violation asymmetries (denoted a_f^d [6]), $a_{D^0 \rightarrow K^+ K^-}^d$ and $a_{D^0 \rightarrow \pi^+ \pi^-}^d$ [13] are also updated.

The decay $D^{*+} \rightarrow D^0 \pi_s^+$ is used to identify the flavor of the D^0 meson from the charge of the low momentum pion (referred to as “the soft pion”), π_s^+ . Thus, we determine $A_{CP}^{D^0 \rightarrow K_S^0 P^0}$ by measuring the asymmetry in the signal yield

$$A_{\text{rec}}^{D^{*+} \rightarrow D^0 \pi_s^+} = \frac{N_{\text{rec}}^{D^{*+} \rightarrow D^0 \pi_s^+} - N_{\text{rec}}^{D^{*-} \rightarrow \bar{D}^0 \pi_s^-}}{N_{\text{rec}}^{D^{*+} \rightarrow D^0 \pi_s^+} + N_{\text{rec}}^{D^{*-} \rightarrow \bar{D}^0 \pi_s^-}}, \quad (2)$$

where N_{rec} is the number of reconstructed decays. The measured asymmetry in Eq. (2) includes two contributions other than A_{CP} . One is the forward-backward asymmetry (A_{FB}) due to $\gamma^* - Z^0$ interference in $e^+e^- \rightarrow c\bar{c}$ and the other is a detection efficiency asymmetry between positively and negatively charged soft pions ($A_{\epsilon}^{\pi^+}$). Since we reconstruct the K_S^0 with $\pi^+\pi^-$ combinations and P^0 with the $\gamma\gamma$ or $\gamma\gamma\pi^+\pi^-$ final states, asymmetries in K_S^0 and P^0 detection cancel out. Equation (2) then can be simplified to give

$$A_{\text{rec}}^{D^{*+} \rightarrow D^0 \pi_s^+} = A_{CP}^{D^0 \rightarrow K_S^0 P^0} + A_{FB}^{D^{*+}}(\cos \theta_{D^{*+}}^{\text{CMS}}) + A_{\epsilon}^{\pi^+}(p_{T\pi_s^+}^{\text{lab}}, \cos \theta_{\pi_s^+}^{\text{lab}}) \quad (3)$$

by neglecting the terms involving the product of asymmetries, where A_{CP} is independent of all kinematic variables, $A_{FB}^{D^{*+}}$ is an odd function of the cosine of the polar angle of D^{*+} in the center-of-mass system (CMS), and $A_{\epsilon}^{\pi^+}$ depends on transverse momentum and polar angle of π_s^+ in the laboratory frame, while it is uniform in azimuthal angle. To correct for $A_{\epsilon}^{\pi^+}$ we use the decays $D^0 \rightarrow K^-\pi^+$ (referred to as untagged) and $D^{*+} \rightarrow D^0\pi_s^+ \rightarrow K^-\pi^+\pi_s^+$ (referred to as tagged), and assumes the same A_{FB} for D^{*+} and D^0 mesons. By subtracting the measured asymmetries in these two decay modes, $A_{\text{rec}}^{\text{untagged}}$ and $A_{\text{rec}}^{\text{tagged}}$, we directly measure the $A_{\epsilon}^{\pi^+}$ correction factor [13, 14]. With $A_{\text{rec}}^{D^{*+} \rightarrow D^0 \pi_s^+}$ corrected for $A_{\epsilon}^{\pi^+}$ (denoted $A_{\text{rec,corr}}^{D^{*+} \rightarrow D^0 \pi_s^+}$ below),

$$A_{\text{rec,corr}}^{D^{*+} \rightarrow D^0 \pi_s^+} = A_{CP}^{D^0 \rightarrow K_S^0 P^0} + A_{FB}^{D^{*+}}(\cos \theta_{D^{*+}}^{\text{CMS}}), \quad (4)$$

we extract A_{CP} and A_{FB} using

$$A_{CP}^{D^0 \rightarrow K_S^0 P^0} = [A_{\text{rec,corr}}^{D^{*+} \rightarrow D^0 \pi_s^+}(\cos \theta_{D^{*+}}^{\text{CMS}}) + A_{\text{rec,corr}}^{D^{*+} \rightarrow D^0 \pi_s^+}(-\cos \theta_{D^{*+}}^{\text{CMS}})]/2, \quad (5a)$$

$$A_{FB}^{D^{*+}} = [A_{\text{rec,corr}}^{D^{*+} \rightarrow D^0 \pi_s^+}(\cos \theta_{D^{*+}}^{\text{CMS}}) - A_{\text{rec,corr}}^{D^{*+} \rightarrow D^0 \pi_s^+}(-\cos \theta_{D^{*+}}^{\text{CMS}})]/2. \quad (5b)$$

The data used in this analysis were recorded at or near the $\Upsilon(4S)$ resonance with the Belle detector [15] at the e^+e^- asymmetric-energy collider KEKB [16]. The sample corresponds to an integrated luminosity of 791 fb^{-1} .

We apply the same charged track selection criteria that were used in Ref. [17]. For soft pions we do not require associated hits in the silicon vertex detector, either in the z or radial directions [18]. Charged kaons and pions are identified by requiring the ratio of particle identification likelihoods [17] to be greater or less than 0.6, respectively. K_S^0 candidates are reconstructed from pairs of oppositely charged tracks that have an invariant mass

within $\pm 9 \text{ MeV}/c^2$ of the nominal K_S^0 mass [9, 17]. Candidate π^0 and η mesons are reconstructed from $\gamma\gamma$ pairs where the minimum energy of each γ is required to be 60 MeV for the barrel and 100 MeV for the forward region of the calorimeter [19]. We require the $\gamma\gamma$ invariant mass to be between 0.11 and 0.16 GeV/c^2 for π^0 candidates and between 0.50 and 0.58 GeV/c^2 for η candidates. The momentum of $\gamma\gamma$ pairs is required to be greater than 0.5 GeV/c for both π^0 and η selections. In order to remove a significant π^0 photon background contribution under the η signal peak, we combine individual γ candidates from $\eta \rightarrow \gamma\gamma$ with any other detected γ in the event. If the $\gamma\gamma$ pair invariant mass is in the π^0 mass window, the γ is rejected. Further reduction of the π^0 contribution under the η signal is achieved by requiring the energy balance of the $\gamma\gamma$ in the η decay to be less than 0.8, where the energy balance is the ratio of the difference and the sum of two γ energies. Candidate η' mesons are reconstructed in the $\eta\pi^+\pi^-$ decay channel. To improve the η' mass resolution, the four-momentum of the η is recalculated with a nominal η mass [9] constraint. The same minimum γ energy requirement used for the $D^0 \rightarrow K_S^0\eta$ selection is imposed in the η' reconstruction. The π^0 veto, however, is not applied since it is found to be unnecessary once the invariant mass of the $\eta\pi^+\pi^-$ candidates is required to be between 0.945 and 0.970 GeV/c^2 and the D^0 mass selection requirement, which is described below, is applied.

The four-momentum of the P^0 is recalculated from a kinematic fit to its nominal mass [9] and combined with a K_S^0 to form a D^0 candidate. D^{*+} candidates are reconstructed using a soft pion and a D^0 candidate with mass in the $[1.75, 1.95] \text{ GeV}/c^2$ ($K_S^0\pi^0$), $[1.82, 1.90] \text{ GeV}/c^2$ ($K_S^0\eta$), or $[1.84, 1.89] \text{ GeV}/c^2$ ($K_S^0\eta'$) interval which depends on the mass resolution. To remove D^{*+} mesons produced in B decays, the D^{*+} momentum in the CMS is required to be greater than 2.5 GeV/c . All selections are chosen to maximize N_S/σ_{N_S} and to minimize the peaking backgrounds, where N_S is the signal yield from the fit and σ_{N_S} is the uncertainty in N_S . After applying all of the selections described above, the $D^0 \rightarrow \pi^+\pi^-\pi^0$ contribution to $D^0 \rightarrow K_S^0\pi^0$ and the $D^0 \rightarrow K_S^0\pi^0$ contribution to $D^0 \rightarrow K_S^0\eta$ are found to be negligible in simulation studies. Figure 1 shows data distributions of the mass difference $M(D^*) - M(D)$ for all the decay modes.

All mass difference signals are parameterized as a sum of a Gaussian and a bifurcated Gaussian distributions with a common mean. The background is parameterized by the form $(x - m_{\pi^+})^\alpha e^{-\beta(x - m_{\pi^+})}$, where α and β are free parameters, m_{π^+} is the charged pion mass [9] and x is the mass difference. The asymmetry and the sum of the D^{*+} and D^{*-} yields are directly obtained from a simultaneous fit to the D^{*+} and D^{*-} candidate distributions. The common parameters in the simultaneous fit are the mean of the Gaussian, the widths of the Gaussian and the bifurcated Gaussian, and the ratio of the Gaussian and the bifurcated Gaussian amplitudes, which are

the same for the $M(D^*) - M(D)$ distributions in different $K_S^0 P^0$ final states and in the slightly different phase spaces of the individual $K_S^0 P^0$ modes. Table I lists the results of the fits.




FIG. 1: Distributions of the mass difference $M(D^*) - M(D)$ for the studied decay modes. Left plots show the mass difference between D^{*+} and D^0 and right plots show that between D^{*-} and $\bar{D}^0$. Top plots are for the $K_S^0 \pi^0$, middle plots for the $K_S^0 \eta$, and bottom plots for the $K_S^0 \eta'$ final states. Points with error bars are the data and the histograms show the results of the parameterizations of the data. Hatched areas are the background contributions.

TABLE I: The sum (N_S) and the asymmetry (A_{rec} in Eq. (2)) of D^{*+} and D^{*-} yields from the fits. The uncertainties are statistical only.

	N_S	A_{rec} (%)
$D^{*+} \rightarrow D^0 \pi_s^+ \rightarrow K_S^0 \pi^0 \pi_s^+$	326303 ± 679	$+0.19 \pm 0.19$
$D^{*+} \rightarrow D^0 \pi_s^+ \rightarrow K_S^0 \eta \pi_s^+$	45831 ± 283	$+1.00 \pm 0.51$
$D^{*+} \rightarrow D^0 \pi_s^+ \rightarrow K_S^0 \eta' \pi_s^+$	26899 ± 211	$+1.47 \pm 0.67$

In order to obtain $A_{\epsilon}^{\pi_s^+}$ we first extract $A_{\text{rec}}^{\text{untagged}}$ using simultaneous fits analogous to those used for the signal modes, but instead of the $M(D^*) - M(D)$ distribution we fit to the $M(D)$ distribution using a similar parameterization. The values of $A_{\text{rec}}^{\text{untagged}}$ are evaluated in bins of transverse momentum ($p_T^{\text{lab}}_{D^0}$) and polar angle ($\cos \theta_{D^0}^{\text{lab}}$) of untagged $D^0 \rightarrow K^- \pi^+$ candidates in the laboratory frame. The p_T and polar angle variables are only weakly correlated. Each tagged $D^* \rightarrow D \pi_s \rightarrow K \pi \pi_s$ candidate is then weighted with $1 - A_{\text{rec}}^{\text{untagged}}$ for D^{*+} and $1 + A_{\text{rec}}^{\text{untagged}}$ for D^{*-} . Details of the weighting procedure are described in Ref. [13]. After this the remaining

asymmetry in the tagged decay sample is $A_{\epsilon}^{\pi_s^+}$, which is obtained from the simultaneous fits to the weighted $M(D^*) - M(D)$ distributions with the same parameterization used in the signal modes, now for bins of $p_T^{\text{lab}}_{\pi_s^+}$ and $\cos \theta_{\pi_s^+}^{\text{lab}}$.

The dominant sources of uncertainty in the $A_{\epsilon}^{\pi_s^+}$ determination are the statistical uncertainties in the untagged and tagged samples. These are found to be 0.04% and 0.07%, respectively. Other sources of systematic uncertainties are found to be negligible. Thus, we assign a total systematic uncertainty of 0.08% to the $A_{\epsilon}^{\pi_s^+}$ determination, obtained by adding the two contributions in quadrature.

The data samples shown in Fig. 1 are divided into bins of p_T^{lab} and $\cos \theta^{\text{lab}}$ of the π_s^+ . The $A_{\epsilon}^{\pi_s^+}$ correction is applied by weighting each D^{*+} event with $1 - A_{\epsilon}^{\pi_s^+}$ and each D^{*-} event with $1 + A_{\epsilon}^{\pi_s^+}$. The weighted mass difference distributions in bins of the D^{*+} polar angle in the CMS are fitted simultaneously to obtain the corrected asymmetry. We fit for the linear component in $\cos \theta_{D^{*+}}^{\text{CMS}}$ to determine A_{FB} while the A_{CP} component is uniform in $\cos \theta_{D^{*+}}^{\text{CMS}}$. Figure 2 shows $A_{CP}^{D^0 \rightarrow K_S^0 P^0}$ and




FIG. 2: Measured A_{CP} (left) and A_{FB} (right) values as a function of $|\cos \theta_{D^{*+}}^{\text{CMS}}|$. Top plots are for $K_S^0 \pi^0$, middle plots for $K_S^0 \eta$, and bottom plots for $K_S^0 \eta'$ final states. The dashed curves show the leading-order prediction for A_{FB}^{c} .

$A_{FB}^{D^{*+}}$ as a function of $|\cos \theta_{D^{*+}}^{\text{CMS}}|$. From a weighted average over the $|\cos \theta_{D^{*+}}^{\text{CMS}}|$ bins, we obtain $A_{CP}^{D^0 \rightarrow K_S^0 \pi^0} = (-0.28 \pm 0.19)\%$, $A_{CP}^{D^0 \rightarrow K_S^0 \eta} = (+0.54 \pm 0.51)\%$, and $A_{CP}^{D^0 \rightarrow K_S^0 \eta'} = (+0.98 \pm 0.67)\%$ where the uncertainties are statistical only. The $\chi^2/\text{d.o.f}$ with respect to the

average over the $|\cos\theta_{D^{*+}}^{\text{CMS}}|$ bins is 5.1/4 ($K_S^0\pi^0$), 3.0/3 ($K_S^0\eta$), or 5.3/3 ($K_S^0\eta'$). The observed A_{FB} values decrease with $\cos\theta_{D^{*+}}^{\text{CMS}}$ as expected from the leading-order prediction [20]. The observed deviations from the prediction are expected due to higher order corrections. Similar A_{FB} 's were found in previous measurements [13, 14, 21]. The results are validated with toy pseudo-experiments and full detector simulation Monte Carlo events. We found no systematic deviations from the input values.

We consider several sources of systematic uncertainty. The uncertainty due to the limited size of the tagged and untagged samples was discussed above. To estimate the systematic uncertainty due to the choice of fitting method and parameters, we vary the histogram binnings, fitting intervals, and signal and background parameterizations. We also consider the systematic uncertainties due to the choice of $\cos\theta_{D^{*+}}^{\text{CMS}}$ binning. Finally, we include possible effects due to the differences in interactions of K^0 and $\bar{K}^0$ mesons with the material of the detector as explained in Ref. [21], and assign a systematic uncertainty of 0.06% due to this effect. Table II summarizes the components of the systematic uncertainties. The larger uncertainties in $A_{CP}^{D^0 \rightarrow K_S^0\eta^{(\prime)}}$ due to the choice of fitting method are a consequence of smaller statistics of these samples.

TABLE II: Summary of systematic uncertainties in A_{CP} .

Source	$K_S^0\pi^0$ (%)	$K_S^0\eta$ (%)	$K_S^0\eta'$ (%)
$A_{\epsilon_s^+}$ determination	0.08	0.08	0.08
Fitting	0.02	0.12	0.10
$\cos\theta_{D^{*+}}^{\text{CMS}}$ binning	<0.01	0.01	0.03
$K^0/\bar{K}^0$ -material effects	0.06	0.06	0.06
Total	0.10	0.16	0.14

From the total uncertainties shown in Table II, we obtain $A_{CP}^{D^0 \rightarrow K_S^0\pi^0} = (-0.28 \pm 0.19 \pm 0.10)\%$, $A_{CP}^{D^0 \rightarrow K_S^0\eta} = (+0.54 \pm 0.51 \pm 0.16)\%$ and $A_{CP}^{D^0 \rightarrow K_S^0\eta'} = (+0.98 \pm 0.67 \pm 0.14)\%$ where the first uncertainties are statistical and the second are systematic. Table III summarizes the results, current world average [9], and $A_{CP}^{\bar{K}^0}$.

TABLE III: Summary of the A_{CP} measurements. The first uncertainties in the second column are statistical and the second are systematic. The third column shows the world average of A_{CP} and the fourth $A_{CP}^{\bar{K}^0}$. $A_{CP}^{D^0 \rightarrow K_S^0\eta^{(\prime)}}$ are the first measurements, hence no world average of A_{CP} is given in the third column.

	Belle (%)	Ref. [9] (%)	$A_{CP}^{\bar{K}^0}$ (%)
$A_{CP}^{D^0 \rightarrow K_S^0\pi^0}$	$-0.28 \pm 0.19 \pm 0.10$	$+0.1 \pm 1.3$	-0.332 ± 0.006
$A_{CP}^{D^0 \rightarrow K_S^0\eta}$	$+0.54 \pm 0.51 \pm 0.16$	—	-0.332 ± 0.006
$A_{CP}^{D^0 \rightarrow K_S^0\eta'}$	$+0.98 \pm 0.67 \pm 0.14$	—	-0.332 ± 0.006

Besides the A_{CP} measurements listed in Table III,

we test the universality of a^{ind} assuming negligible new CP violating effects in D^0 decays to the $K_S^0\pi^0$ final state as discussed in Ref. [6]. By subtracting $A_{CP}^{\bar{K}^0}$ from $A_{CP}^{D^0 \rightarrow K_S^0\pi^0}$, we obtain $a^{\text{ind}} = (+0.05 \pm 0.19 \pm 0.10)\%$, which is consistent with $-A_F = (-0.01 \pm 0.30 \pm 0.15)\%$ obtained in Ref. [2]. This is the first experimental test of a^{ind} in D^0 decays with a sensitivity near 0.3%. By averaging the two independent values we obtain $a^{\text{ind}} = (+0.03 \pm 0.18)\%$, where the uncertainty includes the statistical and systematic errors, and represents the most precise value of a^{ind} from a single-experiment currently. Using the average a^{ind} , we also update the values of $a_{D^0 \rightarrow K^+K^-}^d$ and $a_{D^0 \rightarrow \pi^+\pi^-}^d$ from Ref. [13], which are $(-0.46 \pm 0.37)\%$ and $(+0.40 \pm 0.56)\%$ [22], respectively. The errors include all the uncertainties of input measurements.

In summary, we report a search for CP violation in the decays $D^0 \rightarrow K_S^0 P^0$ using a data sample with an integrated luminosity of 791 fb^{-1} collected with the Belle detector. We observe no evidence for CP violation. The measurement in the decay $D^0 \rightarrow K_S^0\pi^0$ is the most precise measurement of any CP asymmetry in the charmed particle sector to date. We also report the first measurements of CP asymmetries in the decays $D^0 \rightarrow K_S^0\eta$ and $D^0 \rightarrow K_S^0\eta'$. Our results are consistent with the SM and can be used to place the most stringent constraints on NP models arising from the measurements of CP violation in the charm sector at present.

We thank the KEKB group for excellent operation of the accelerator, the KEK cryogenics group for efficient solenoid operations, and the KEK computer group and the NII for valuable computing and SINET3 network support. We acknowledge support from MEXT, JSPS and Nagoya's TLP RC (Japan); ARC and DIISR (Australia); NSFC (China); MSMT (Czechia); DST (India); MEST, NRF, NSDC of KISTI, and WCU (Korea); MNiSW (Poland); MES and RFAAE (Russia); ARRS (Slovenia); SNSF (Switzerland); NSC and MOE (Taiwan); and DOE (USA). B. R. Ko acknowledges support by NRF Grant No. 2010-0021279 and E. Won acknowledges support by NRF Grant No. 2009-0071072.

-
- [1] B. Aubert *et al.* (Babar Collab.), Phys. Rev. Lett. **98**, 211802 (2007).
 - [2] M. Starič *et al.* (Belle Collab.), Phys. Rev. Lett. **98**, 211803 (2007).
 - [3] T. Aaltonen *et al.* (CDF Collab.), Phys. Rev. Lett. **100**, 121802 (2008).
 - [4] D. Asner *et al.* (Heavy Flavor Averaging Group), [arXiv:1010.1589v1\[hep-ex\]](https://arxiv.org/abs/1010.1589v1) and online update at <http://www.slac.stanford.edu/xorg/hfag/>.
 - [5] A. F. Falk, Y. Grossman, Z. Ligeti, and A. A. Petrov, Phys. Rev. D **65**, 054034 (2002); A. F. Falk, Y. Grossman, Z. Ligeti, Y. Nir, and A. A. Petrov, Phys. Rev. D

- 69**, 114021 (2004).
- [6] Y. Grossman, A. L. Kagan, and Y. Nir, Phys. Rev. D **75**, 036008 (2007).
- [7] F. Buccella, M. Lusignoli, G. Miele, A. Pugliese, and P. Santorelli, Phys. Rev. D **51**, 3478 (1995).
- [8] Throughout this Letter the charge-conjugate decay mode is also implied unless stated otherwise.
- [9] K. Nakamura *et al.* (Particle Data Group), J. Phys. G **37**, 075021 (2010).
- [10] We use $A_{CP}^{\bar{K}^0} = \frac{\Gamma(\bar{K}^0 \rightarrow \bar{f}) - \Gamma(K^0 \rightarrow f)}{\Gamma(\bar{K}^0 \rightarrow \bar{f}) + \Gamma(K^0 \rightarrow f)}$. Hence, $A_{CP}^{\bar{K}^0} = -A_L$ where A_L is the symbol for the asymmetry used in Ref. [9].
- [11] M. Kobayashi and T. Maskawa, Prog. Theor. Phys., **49**, 652 (1973).
- [12] I. I. Bigi and H. Yamamoto, Phys. Lett. B **349**, 363 (1995).
- [13] M. Starič *et al.* (Belle Collab.), Phys. Lett. B **670**, 190 (2008).
- [14] B. Aubert *et al.* (Babar Collab.), Phys. Rev. Lett. **100**, 061803 (2008).
- [15] A. Abashian *et al.* (Belle Collab.), Nucl. Instr. and Meth. A **479**, 117 (2002).
- [16] S. Kurokawa and E. Kikutani, Nucl. Instr. and Meth. A **499**, 1 (2003), and other papers included in this volume.
- [17] E. Won *et al.* (Belle Collab.), Phys. Rev. D **80**, 111101(R) (2009).
- [18] Z. Natkaniec *et al.* (Belle SVD2 Group), Nucl. Instr. and Meth. A **560**, 1 (2006); Y. Ushiroda (Belle SVD2 Group), Nucl. Instr. and Meth. A **511**, 6 (2003).
- [19] H. Ikeda *et al.*, Nucl. Instr. and Meth. A **441**, 401 (2000).
- [20] The leading-order prediction for A_{FB}^{cc} at $\sqrt{s}=10.6$ GeV is about $-0.029 \cos \theta^{\text{CMS}} / (1 + \cos^2 \theta^{\text{CMS}})$. See for example O. Nachtmann, Elementary Particle Physics, Springer-Verlag (1989).
- [21] B. R. Ko *et al.* (Belle Collab.), Phys. Rev. Lett. **104**, 181602 (2010).
- [22] The systematic uncertainty due to the $A_{\epsilon_s}^{\pi_s^+}$ determination in the two measurements are only partially correlated due to the use of slightly different data set and different tracking algorithms. The difference in the results when treating the systematic uncertainties as completely correlated or completely uncorrelated is negligible. To be conservative, we quote the result treating the systematic uncertainties as uncorrelated.